

DERMATOMYCOSIS VACCINE

Related Applications

The present application is a continuation of United States application Serial No. 09/256,915,
5 filed February 24, 1999, which is a continuation of United States application Serial No.
08/568,063, filed December 6, 1995, now abandoned, which is a continuation of United
States application Serial No. 08/281,380, filed July 26, 1994, now abandoned, which is a
continuation of United States application Serial No. 08/081,299, filed August 11, 1993, now
abandoned, which claims priority under 35 USC § 119 to Russian Federation application
10 Serial No. 50068611307308, filed October 21, 1991; and claims priority benefit of all the
above-listed applications.

Background

This invention relates to the preparation of vaccines and their use in preparing compositions
15 for specifically preventing and treating dermatomycosis.

Dermatomycoses in animals are anthroponozoonotic diseases of the skin and related tissue.
Clinical symptoms are characterized by loss of hair in the affected area, hyperemia, scaling
and asbestos-like scabs. Inflammation is often accompanied by suppuration.
20 Dermatomycoses are often also characterized by localized infection of the skin.

Dermatomycoses in animals carry a substantial socioeconomic impact. Diseased animals
required prolonged treatment and can spread infection to both animals and humans.

25 Up till now, dermatomycoses have been treated using various types of medication applied
locally to affected areas of the skin. These included the ointments YaM, Yuglon (I) and a
number of other ointments, liniments, solutions and other substances containing fungicides
and fungistatic agents.

30 The disadvantages of such treatments were:

they required the adoption of quarantine measures and disinfection of areas, etc.;
animals were kept (rearing pens, vivaria, farms, zoos, circuses, etc.);

- 5

10

15

20

25

It has also been noted that the areas where a live vaccine is injected may become a specific focus in which cultures of vaccinal strains may at certain times be produced. Given that some species of domestic animals come into frequent contact with humans, the occurrence of such

Description

This invention now provides universal inactivated vaccines for the specific treatment and prevention of dermatomycosis in animals and corresponding immunogenic fungal strains.

- 5 This aim has been achieved by using the following fungal strains as vaccinal strains:

Trichophyton verrucosum (especially No. VKPGF-931/410), *Trichophyton mentagrophytes* (especially No. VKPGF-930/1032), *Trichophyton equinum* (especially No. VKPGF-929/381), *Trichophyton sarkisovii* (especially No. VKPGF-551/68), *Microsporum canis* (especially No. VKPGF-928/1393), *Microsporum canis* var. *obesum* (especially No. VKPGF-727/1311),
 10 *Microsporum canis* var. *distortum* (especially No. VKPGF-728/120), *Microsporum gypseum* (especially No. VKPGF-729/59). Vaccines can be produced by using various combinations of antigenic material from the above strains together with a suitable carrier.

A preferred combination consists of *Trichophyton verrucosum* No. VKPGF-931/410,

- 15 *Trichophyton mentagrophytes* No. VKPGF-930/1032, *Trichophyton equinum* No. VKPGF-929/381, *Microsporum canis* No. VKPGF-928/1393, *Microsporum canis* var. *obesum* No. VKPGF-727/1311, *Microsporum canis* var. *distortum* No. VKPGF-728/120, *Microsporum gypseum* No. VKPGF-729/59, particularly for use in dogs, cats and horses.

- 20 Another preferred combination of vaccine strains consists of *Trichophyton verrucosum* No. VKPGF-931/410, *Trichophyton mentagrophytes* No. VKPGF-930/1032, *Trichophyton sarkisovii* No. VKPGF-551/68, particularly for use in cattle.

- 25 The antigenic material may comprise a single antigen of at least one, and more particularly of all of the above-mentioned dermatophytes or from a plurality of antigens, provided that a sufficient immune response is stimulated to give resistance to a dermatophyte infection.

Antigenic material for such a purpose can be prepared using methods known from the prior art, e.g., homogenizing the above-mentioned dermatophytes or parts thereof, fractionation of dermatophyte preparations, production of antigenic dermatophyte material by recombinant DNA technology, etc. It is preferably to use a homogenized culture material having 10 to 12

Suitable physiologically acceptable carriers for administering the vaccines are known from the prior art and may include buffers, gels, microparticles, implantable solids, solutions and other adjuvants.

- 5 To kill off the dermatophytes it is possible to use thiomersal ($C_9H_9O_2SNaHg$), formaldehyde or 2-propiolactone.

In order to prepare a vaccine the following procedure may be used, for example:

- 10 Cultures of the strains are homogenized in an aqueous solution containing 0.2 to 2.0% fermented, hydrolyzed muscle protein (FGM-s), 5 to 12% glucose and 0.1 to 1.2% yeast extract. The concentration of the microconidia is adjusted to 40 to 120 million per milliliter and after 1 to 2 days the mixture is inactivated, *e.g.*, with thiomersal ($C_9H_9O_2SNaHg$) in the
15 resulting suspension is packaged and is ready for use in animals.

The preparation of the vaccines, the dosage to be given and the method of administration for prevention and therapeutic treatment are explained in Examples 1 to 3.

- 20 The invention now makes it possible to prepare an inactivated vaccine that reduces the probability of reinfection and also implants a high degree of immunity. Unlike the known vaccines, the vaccine according to the invention in practice gives immunity to all important causes of dermatomycosis in animals.

- 25 Briefly, the vaccine according to the invention offers the following advantages:
- in many species of disease-prone animals it establishes immunity after intramuscular injection,
 - it grants immunity against almost all causes of dermatomycosis in animals,
 - it has stable immunogenic properties.

It is easy to prepare

shows no side reactions in animals.

The vaccine has been successfully tested on over 500 animals of different species, predominantly in affected regions.

5 The strains used to produce the vaccine have been deposited at the "All-Union Collection of Pathogenic Fungi within the USSR, Ministry of Health Centre for Deep Mycoses" in Leningrad and at the "DSM – Deutsche Sammlung von Mikroorganismen und Zellkulturen", Mascheroder Weg 1B, W-3300 Braunschweig, Germany.

10 The following microorganisms were deposited with the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSM) on October 1, 1992 and received the following accession numbers:

- *T. verrucosum* VKPGF-931/410 received accession No. DSM 7277
- *T. mentagrophytes* VKPGF-930/1032 received accession No. DSM 7279
- 15 - *T. equinum* VKPGF-929/381 received accession No. DSM 7276
- *T. sarkisovii* VKPGF-551/68 received accession No. DSM 7278
- *M. canis* VKPGF-928/1393 received accession No. DSM 7281
- *M. canis var. obesum* VKPGF-727/1311 received accession No. DSM 7280
- *M. canis var. distortum* VKPGF-728/120 received accession No. DSM 7275
- 20 - *M. gypseum* VKPGF-729/59 received accession No. DSM 7274.

DSM is located at Macheroder Weg 1B, W-3300 Braunschweig, Germany. "T" is an abbreviation for *Trichophyton* and "M" is an abbreviation for *Microsporum*.

25 Their characteristics are set out below:

TRICHOPHYTON VERRUCOSUM, No. VKPGF-931/410

The strain was deposited at the DSM – Deutsche Sammlung von Mikroorganismen und Zellkulturen", Mascheroder Weg 1B, W-3300 Braunschweig, Germany.

The epizootic strain No. 931/410, which was identified on a deer in 1988, the strain was identified using the Rebell-Taplin key (Rebell, G., Taplin, D.: Dermatophytes, their

recognition and identification, 1978) and according to Kashkin, P.N. *et al.* (Opredelitel patogennykh, toksigenykh vrednykh dlya cheloveka gribov, 1979).

The biological properties of the strain are described in Table 1.

5

Strain No. VKPGF-931/410 differs from the epizootic strain in its faster growth in nutrient medium, the enormous production of microconidia, lower virulence and the absence of any reaction with its antigens.

10 TRICHOPHYTON MENTAGROPHYTES No. VKPGF-930/1032

The strain was deposited at the DSM – Deutsche Sammlung von Mikroorganismen und Zellkulturen”, Mascheroder Weg 1B, W-3300 Braunschweig, Germany.

15 The strain was obtained by directed selection based on spore production and attenuation of the epizootic Strain No. 1032, which was found on a horse in 1985. The strain was identified as described above Rebel, Taplin, *loc. cit.* and Kashkin, *loc. cit.*). The biological properties are described in Table 2.

20 Strain No. VKPGF-930/1032 differs from the epizootic strain by its faster growth in nutrient medium, the enormous production of microconidia, its lower virulence and the absence of any reaction with its antigens.

TRICHOPHYTON EQUINUM No. VKPGF-929/381

25 The strain was deposited at the DSM – Deutsche Sammlung von Mikroorganismen und Zellkulturen”, Mascheroder Weg 1B, W-3300 Braunschweig, Germany.

The strain was obtained by directed selection based on spore production and attenuation of the epizootic Strain No. 381 which was found on a horse in 1986. The strain was identified as describe above Rebel, Taplin, *loc. cit.* and Kashkin, *loc. cit.*). The biological properties are
30 described in Table 3.

Strain No. VKPGF-929/381 differs from the epizootic strain by its faster growth in nutrient medium, lower virulence and the absence of any reaction with its antigens.

MICROSPORUM CANIS No. VKPGF-928/1393

The strain was deposited at the DSM – Deutsche Sammlung von Mikroorganismen und Zellkulturen”, Mascheroder Weg 1B, W-3300 Braunschweig, Germany.

5

The strain was obtained by directed selection based on spore production and attenuation of the epizootic Strain No. 1393 which was found on a cat in 1988. The strain was identified as describe above Rebel, Taplin, *loc. cit.* and Kashkin, *loc. cit.*). The biological properties are described in Table 4.

10

Strain No. VKPGF-928/1393 differs from the epizootic strain by its faster growth in nutrient medium, its enormous capacity to carry spores, lower virulence and the absence of any reaction with its antigens.

15 MICROSPORUM CANIS VAR. OBESUM No. VKPGF-727/1311

The strain was deposited at the DSM – Deutsche Sammlung von Mikroorganismen und Zellkulturen”, Mascheroder Weg 1B, W-3300 Braunschweig, Germany.

20

The strain was obtained by directed selection based on spore production and attenuation of the epizootic Strain No. 1311 which was found on a tiger in 1986. The strain was identified as describe above Rebel, Taplin, *loc. cit.* and Kashkin, *loc. cit.*). The biological properties are described in Table 5.

25

Strain No. VKPGF-727/1311 differs from the epizootic strain by its faster growth in nutrient medium, its enormous capacity to carry spores, lower virulence and the absence of any reaction with its antigens.

MICROSPORUM CANIS VAR. DISTORTUM No. VKPGF-728/120

30

The strain was deposited at the DSM – Deutsche Sammlung von Mikroorganismen und Zellkulturen”, Mascheroder Weg 1B, W-3300 Braunschweig, Germany.

The strain was obtained by directed selection based on spore production and attenuation of the epizootic Strain No. 120 which was found on a black panther in 1987. The strain was

identified as describe above Rebel, Taplin, *loc. cit.* and Kashkin, *loc. cit.*). The biological properties are described in Table 6.

5 Strain No. VKPGF-728/120 differs from the epizootic strain by its faster growth in nutrient medium, its enormous production of microconidia, its lower virulence and the absence of any reaction with its antigens.

MICROSPORUM GYPSEUM No. VKPGF-729/59

10 The strain was deposited at the DSM – Deutsche Sammlung von Mikroorganismen und Zellkulturen”, Mascheroder Weg 1B, W-3300 Braunschweig, Germany.

The strain was obtained by directed selection based on spore production and attenuation of the epizootic Strain No. 59 which was found on a horse in 1985. The strain was identified as describe above Rebel, Taplin, *loc. cit.* and Kashkin, *loc. cit.*). The biological properties are
15 described in Table 7.

Strain No. VKPGF-729/59 differs from the epizootic strain by its faster growth in nutrient medium, the enormous production of microconidia, the lower virulence and the absence of any reaction with its antigens.

No.	Name and characteristics of strain	Strain No. VKPGF-931/410	Epizootic Strain No. 410
1	General characteristics	Mature 10-15 day single-spore colony in agar/wort; white, velvety, convex, narrow growing margin undersurface colorless, colony diameter 10-15 mm	Mature 25-30 day colony in agar/wort; cream, leathery/velvety, folded, undersurface colorless, colony diameter 9-13 mm
2	Morphological characteristics	Mature 10-15 day culture with septate branching hyphae 1-3 µm wide, numerous oval, pyriform microconidia measuring 1.5 to 3 x 3 to 5 µm, no macroconidia	Mature 25-30 day culture with septate branching mycelium 1-3 µm wide, few oval, pyriform, cylindrical microconidia measuring 1 to 3 x 3 to 7 µm, single elongate irregular shape macroconidia with 2-5 septates measuring 3 to 5 x 25 to 30 µm, numerous arthrospores in chains 6-8 µm diameter, chlamydospores 10-12 µm diameter
3	Toxic characteristics	Days after application of a dose of 500-600 cells of fungal matter per cm ² to the scarified rabbit	Dense asbestos-like scabs, possible suppuration
4	Recovery after	19-20 days	25-30 days
5	Response	No observed changes in clinical state	Inflammation at point of injection, edema
6	Response of subcutaneous and intramuscular injection of corpuscular antigens from cultures	1:320 to 1:640 1:400 to 1:1600	1:320 to 1:640 1:400 to 1:1600
7	Response	Establishes immunity	Establishes immunity
8	Response	Establishes immunity	Establishes immunity
9	Response	Establishes immunity	Establishes immunity
10	Response	Establishes immunity	Establishes immunity
11	Response	Establishes immunity	Establishes immunity

Table	Characteristics of strain	Strain No. VKPGF-930/1032	Epizootic Strain No. 1032
Properties	Culture	Mature 10-15 day colony in agar/wort; cream, velvety/powdered, flat with slight flat elevation in center, narrow growing margin, fringed, undersurface light brown, colony diameter 25-30 mm	Mature 25-30 day colony in agar/wort; white, flat, narrow growing margin, undersurface reddish-brown, colony diameter 15-20 mm
Morphology	Colonial characteristics	Septate, branching hyphae 1-3 µm wide, numerous pyriform, oval microconidia measuring 1 to 3 x 2 to 6 µm, no macroconidia	Septate, branching straight and spiral hyphae 1-3 µm wide, round, flattened pyriform microconidia measuring 1 to 3 x 2 to 6 µm, few elongate-oval macroconidia with 2-5 septates, measuring 2 to 6 x 15 to 25 µm
Pathogenicity	Characteristics after application of a dose of 500-600 mg of fungal matter per cm ² to the scarified skin	Necrotic scabs	Dense, asbestos-like scabs
Spore recovery	Recovery after	22-25 days	30-35 days
Reactions	Response to cutaneous and intramuscular injection of corpuscular antigens from cultures	No observed changes in clinical state	Inflammation at point of injection, edema
Antigenic properties	Response after injecting rabbits with corpuscular body titers observed in blood serum	1:320 to 1:640 1:400 to 1:1600	1:320 to 1:640 1:400 to 1:1600
Immunity	Response of a group of rabbits with inactivated cultures (repeated at least 5 times)	Establishes immunity	Establishes immunity

Table	Characteristics of strain	Strain No. VKPGF-929/381	Epizootic Strain No. 381
Properties	Culture	Mature 10-15 day colony in agar/wort; white, velvety/powdery, flat with slight elevation in center, narrow growing margin, fringed, undersurface light brown, colony diameter 15-20 mm	Mature 15 day colony in agar/wort; white, velvety, slightly creased center, narrow growing margin, undersurface reddish-brown, colony diameter 13-15 mm
Morphological characteristics	Colonial characteristics	Septate, branching hyphae 1-3 µm wide, numerous oval pyriform microconidia measuring 2 to 3 x 3 to 6 µm, no macroconidia	Septate, branching hyphae with coil end, 1-4 µm wide, few oval, pyriform microconidia measuring 2 to 3 x 3 to 7 µm, lobar macroconidia measuring 4 to 7 x 15 to 25 µm
Pathogenicity	Pathogenicity after application of a dose of 500-600 mg of fungal matter per cm ² to the scarified skin	Necrotic scabs	Asbestos-like scabs
Spore recovery	Spore recovery after	20-22 days	25-30 days
Reaction	Reaction to cutaneous and intramuscular injection of corpuscular antigens from cultures	No observed changes in clinical state	Inflammation at point of injection, edema
Antigenic properties	Antigenic properties after injecting rabbits with corpuscular body titers observed in blood serum	1:320 to 1:640 1:800 to 1:1600	1:320 to 1:640 1:800 to 1:1600
Immunogenic properties	Immunogenic response of a group of rabbits with inactivated cultures (repeated at least 5 times)	Establishes immunity	Establishes immunity

Table			
Proj	Characteristics of strain	Strain No. VKPGF-928/1393	Epizootic Strain No. 1393
Description		Mature 10-15 day colony in agar/wort; white, fluffy, convex, narrow growing margin, arachnoid, undersurface brown, colony diameter 30-35 mm	Mature 15 day colony in agar/wort; greyish-beige, arachnoid, powdery in center, growing margin fringed, undersurface yellowish, colony diameter 20-25 mm
Morphological characteristics		Septate, branching hyphae 1-4 μ m wide, numerous pyriform, cylindrical microconidia, few fusiform macroconidia with 3-11 septates, measuring 10 to 20 x 40 to 75 μ m	Septate, branching hyphae 2 to 6 μ m wide, few pyriform, cylindrical microconidia measuring 1 to 3 x 3 to 7 μ m, numerous fusiform macroconidia with 3-11 septates, measuring 10 to 20 x 45 to 85 μ m
Pathogenicity 9 to 10% mortality after application of a dose of 500-600 thousands of a.c. Spores		Necrotic scabs 20-24 days	Dense, asbestos-like scabs 25-45 days
Reaction Results inactivated	Subcutaneous and intramuscular injection of corpuscular antigens from cultures	No observed changes in clinical state	Edema and inflammation at point of injection
Antigen 20 to 25% antigen By P.P.I. By I.I.	Antigen after injecting rabbits with corpuscular body titers observed in blood serum	1:320 to 1:640 1:400 to 1:1600	1:320 to 1:640 1:400 to 1:1600
Immune response of a group of rabbits with inactivated cultures (repeated at least 5 times)		Establishes immunity	Establishes immunity

Table				
Proj	Characteristics of strain	Strain No. VKPGF-727/1311	Epizootic Strain No. 1311	
Description	if culture	Mature 10-15 day colony in agar/wort; white, fluffy, flat with a denser central dome-like elevation, narrow growing margin, fringed, undersurface colorless with brown center, colony diameter 30-35 mm	Mature 15 day colony in agar/wort; greyish, fasciculate/arachnoid with pieces of cottony white mycelium, growing margin fringed, undersurface brownish, colony diameter 23-28 mm	
Morphological characteristics		Septate, branching hyphae 1-3 μ m wide, numerous pyriform, oval and cylindrical microconidia measuring 1 to 3 x 3 to 7 μ m, few short, elliptical, fusiform, elongate-oval macroconidia, some irregular shapes, less frequently "beaked", with 2-5 septates, measuring 11 to 20 x 25 to 50 μ m	Septate, branching hyphae 1-5 μ m wide, few oval, cylindrical microconidia measuring 1 to 3 x 3 to 8 μ m, numerous elliptical, fusiform, elongate-oval or irregularly-shaped macroconidia with 2-5 septates, measuring 11 to 20 x 25 to 55 μ m	
Pathology	Characteristics after application of a dose of 500-600 μ g of fungal matter per cm^2 to the scarified skin	Thin necrotic scabs	Dense, asbestos-like scabs	
Spore	recovery after	10-25 days	25-30 days	
Reaction	Response to cutaneous and intramuscular injection of corpuscular antigens from cultures	No observed changes in clinical state	Inflammation and edema at point of injection	
Results	Response after injecting rabbits with corpuscular antigens observed in blood serum	1:320 to 1:640 1:800 to 1:1600	1:320 to 1:640 1:800 to 1:1600	
Antigen	Response of a group of rabbits with inactivated antigen cultures (repeated at least 5 times)	Establishes immunity	Establishes immunity	

Table			
Properties	Characteristics of strain	Strain No. VKPGF-728/120	Epizootic Strain No. 120
Description	Culture	Mature 10-15 day colony in agar/wort; cream, velvety/powdery, button-like elevation in center, narrow growing margin, finely-fringed, undersurface light-brown with dark-brown center, colony diameter 25-30 mm	Mature 15 day colony in agar/wort; light-beige, powdery, umbonate, narrow growing margin, undersurface brown, colony diameter 18-20 mm
Morphological characteristics	Morphological characteristics	Septate, branching hyphae 1-3 μ m wide, numerous pyriform, oval, cylindrical microconidia measuring 1 to 3 x 3 to 8 μ m, few irregular deformed macroconidia, distorted or fusiform with 2-9 septates, measuring 8 to 20 x 25 to 70 μ m	Septate, branching hyphae 1-3 μ m wide, few pyriform, oval, cylindrical microconidia measuring 1 to 3 x 3 to 8 μ m, numerous irregular deformed or fusiform macroconidia with 2-9 septates, measuring 8 to 20 x 25 to 80 μ m
Pathogenesis	Pathogenesis	after application of a dose of 500-600 μ g of fungal matter per cm^2 to the scarified skin	
Spore recovery after	Spore recovery after	20-25 days	27-45 days
Reaction	Reaction	No observed changes in clinical state	Inflammation and edema at point of injection
Results of inactivation	Results of inactivation	Thin necrotic scabs	Asbestos-like scabs
Antibody response	Antibody response	1:320 to 1:640 1:800 to 1:1600	1:320 to 1:640 1:800 to 1:1600
Antibody titer	Antibody titer	Establishes immunity	Establishes immunity
Antibody response of a group of rabbits with inactivated cultures (repeated at least 5 times)	Antibody response of a group of rabbits with inactivated cultures (repeated at least 5 times)	Establishes immunity	Establishes immunity

Table			
Prop.	Characteristics of strain	Strain No. VKPGF-729/59	Epizootic Strain No. 59
Description	Culture	Mature 10-15 day colony in agar/wort; white, velvety/fluffy, flat with slight elevation in center of colony, flat growing margin, undersurface brownish, colony diameter 25-30 mm	Mature 15 day colony in agar/wort; cream, velvety/powdery, flat with fluffy white mycelium in center, thin growing margin, undersurface brownish, colony diameter 20-22 mm
Morphological characteristics		Septate, branching hyphae 2-3 μ m wide, numerous oval, pyriform, cylindrical microconidia measuring 2 to 4 x 3 to 7 μ m, no or few macroconidia, elliptical, elongate-oval shape with 2-5 septates, measuring 7 to 15 x 25 to 40 μ m	Septate, branching hyphae 2-5 μ m wide, few oval, pyriform, cylindrical microconidia measuring 2 to 4 x 3 to 7 μ m, numerous elliptical, stretched-oval macroconidia with 2-5 septates, measuring 7 to 15 x 25 to 50 μ m
Pathogenicity	after application of a dose of 500-600 μ g of fungal matter per cm ² to the scarified skin	Thin necrotic scabs	Dense, asbestos-like scabs
Spore recovery after		20-22 days	25-28 days
Reaction	Response to cutaneous and intramuscular injection of corpuscular antigens from cultures	No observed changes in clinical state	Inflammation at point of injection
Antibody response	after injecting rabbits with corpuscular antigens observed in blood serum	1:320 to 1:640 1:400 to 1:1600	1:320 to 1:640 1:400 to 1:1600
Immune response	of a group of rabbits with inactivated cultures (repeated at least 5 times)	Establishes immunity	Establishes immunity

The vaccine may be prepared using the strain *Trichophyton sarkovii*, No. 551/68. It is described for example in USSR Patent No. 1177972 dated 08.08.1985, to which reference is made in its entirety.

- 5 This strain was also deposited at the DSM – Deutsche Sammlung von Mikroorganismen und Zellkulturen, Mascheroder Weg 1B, W-3300 Braunschweig, Germany.

In particular, the invention relates to the following:

- 10 - a dermatomycosis vaccine, characterized in that it contains antigenic material from at least one of the following dermatophytes:
- *Trichophyton verrucosum*, particularly *Trichophyton verrucosum* Strain No. VKPGF-931/410 and/or
 - *Trichophyton mentagrophytes*, particularly *Trichophyton mentagrophytes* Strain
 - 15 No. VKPGF-930/1032 and/or
 - *Trichophyton sarkisovii*, particularly *Trichophyton sarkisovii* Strain No. VKPGF-551/68 and/or
 - *Microsporum canis*, particularly *Microsporum canis* Strain No. VKPGF-928/1393 and/or
 - 20 - *Microsporum canis* var. *obesum*, particularly *Microsporum canis* var. *obesum* Strain No. VKPGF-727/1311 and/or
 - *Microsporum canis* var. *distortum*, particularly *Microsporum canis* var. *distortum* Strain No. VKPGF-728/120 and/or
 - *Microsporum gypseum*, particularly *Microsporum gypseum* Strain No. VKPGF-
 - 25 729/59,

and a physiologically acceptable carrier.

- a dermatomycosis vaccine, particularly as an agent for treating dogs, cats and horses, characterized in that it contains antigenic material from the dermatophyte strains
- 30 *Trichophyton verrucosum* No. VKPGF-931/410, *Trichophyton mentagrophytes* No. VKPGF-930/1032, *Trichophyton sarkisovii* No. VKPGF-551/68, *Microsporum canis* No. VKPGF-928/1393, *Microsporum canis* var. *obesum* No. VKPGF-727/1311, *Microsporum canis* var. *distortum* No. VKPGF-728/120 and *Microsporum gypseum* No. VKPGF-729/59.

VKPGF-728/120, *Microsporum gypseum* No. VKPGF-729/59, together with a physiologically acceptable carrier.

- a dermatomycosis vaccine, more particularly as an agent for treating cattle, characterized in that it contains antigenic material from the dermatophyte strains *Trichophyton verrucosum* No. VKPGF-931/410, *Trichophyton mentagrophytes* No. VKPGF-930/1032, *Trichophyton equinum* No. VKPGF-929/381, *Trichophyton sarkisovii* Strain No. VKPGF-551/68, together with a physiologically acceptable carrier.

- a dermatomycosis vaccine as described above, characterized in that it contains 40 to 120 million, preferably 90 million microconidia,

- a dermatomycosis vaccine as described above, characterized in that it contains thiomersal or formaldehyde or 2-propiolactone as inactivator,

- a dermatomycosis vaccine as described above, characterized in that the physiologically acceptable carrier used is an aqueous solution containing 0.2 to 2.0 percent weight of fermented, hydrolyzed muscle protein, 5 to 12 percent weight glucose and 0.1 to 1.2 percent weight yeast extract,

- the dermatophyte strains:

- *Trichophyton verrucosum* Strain No. VKPGF-931/410,
- *Trichophyton mentagrophytes* Strain No. VKPGF-930/1032,
- *Trichophyton equinum* Strain No. VKPGF-929/381,
- *Microsporum canis* Strain No. VKPGF-928/1393,
- *Microsporum canis* var. *obesum* Strain No. VKPGF-727/1311,
- *Microsporum canis* var. *distortum* Strain No. VKPGF-728/120, and
- *Microsporum gypseum* Strain No. VKPGF-729/59.

- a process for preparing a vaccine, characterized in that

- *Trichophyton mentagrophytes* Strain No. VKPGF-930/1032,
- *Trichophyton mentagrophytes* Strain No. VKPGF-930/1032,

- *Trichophyton sarkovii* Strain No. VKPGF-551/68,
 - *Microsporum canis* Strain No. VKPGF-928/1393,
 - *Microsporum canis* var. *obesum* Strain No. VKPGF-727/1311,
 - *Microsporum canis* var. *distortum* Strain No. VKPGF-728/120,
 - 5 - *Microsporum gypseum* Strain No. VKPGF-729/59, and
- b. the antigenic material is mixed with a physiologically acceptable carrier.
- a process as described above, characterized in that an agent, particularly thiomersal, formaldehyde or 2-propiolactone is added to inactivate the dermatophytes.

10

The invention is illustrated by means of the Examples that follow.

Examples

15 Example 1

To produce 1 liter of vaccine, cultures are taken of the strains VKPGF-931/410, 930/1032, 929/381, 551/68, 928/1393, 727/1311, 728/120, and 729/59 and grown in agar/wort at 26°C for 15 days. Each culture is grown in 8 mattress flasks. The fungal mass is then lifted off, homogenized, placed in 200 ml of solution and added to each mixer. The solution used is an aqueous solution containing 1% fermented hydrolyzed muscle protein, 10% glucose and 1% yeast extract. The concentration of microconidia is brought to 90 million per ml of homogenate. After 2 days, 125 ml of each culture in suspension is taken and mixed in a single container. The vaccine may be prepared by mixing together various combinations of the given strains.

25

To inactivate the homogenate mixture, thiomersal is added directly to the cell suspension in the ration 1:20,000. 50 mg of thiomersal is added for every liter of homogenate. The cell mixture is allowed to stand at room temperature for 2 days.

30 The resulting vaccine is bottled, checked for sterility, safety and immunogenic properties in

Vaccine produced in this manner was used to immunize animals.

For prophylactic and treatment purposes the vaccine was used in the following doses (see Table 8):

Animal family	Age	Site of injection	Dosage (ml)	
			Prophylactic	Treatment
Canidae medium/large	1-6 months	Gluteal muscles	2-5	3-6
	6 months +	Gluteal muscles	3-7	4-10
Feline cats	1-5 months	Gluteal muscles	1-1.5	1-1.5
	5 months +	Gluteal muscles	1-2	1-2
Canidae	1-12 months	Gluteal muscles	1-3	3-5
	12 months +	Gluteal muscles	3-5	5-6
Canidae	1-10 months	Gluteal muscles	0.3-0.5	0.5
	10 months +	Gluteal muscles	0.3-0.5	0.5-1.0
Canidae	1-12 months	Gluteal muscles	0.3-0.5	0.5
	12 months +	Gluteal muscles	0.5-1.0	0.5-1.0
Canidae	1-12 months	Gluteal muscles	1-3	1-3
	12 months	Gluteal muscles	3-5	5-6
Canidae	1-10 months	Gluteal and shoulder muscles	0.3-0.5	0.5-1.0
	10 months +	Neck area	0.3-1.0	0.5-1.0
Canidae	3-12 months	Neck area	0.3-0.5	0.5-1.0
	12 months +	Neck area	0.5	0.5-1.0
Canidae	1-6 months	Shoulder and neck area	3-5	5-10
	6 months +	Neck area	5-8	7-10
Canidae	1-12 months	Neck area	3-5	5-10
	12 months	Neck area	5-8	7-10

Example 2

The vaccine produced by the method described in Example 1 was tested on laboratory animals and various other animals for effectiveness in the prevention and treatment of disease. The results are given in Table 9.

5

Example 3

The vaccine produced by the method described in Example 1 was also used to treat animals suffering from dermatophytiae. The results are given in Table 10.

Table 9			
Type of animals	Number	Dosage (cm ³)	Effectiveness
Rabbits	10	1.0	No symptoms of disease after infection with virulent cultures of the fungi <i>T. mentagrophytes</i> , <i>T. verrucosum</i> , <i>T. equinum</i> , <i>M. canis</i> , <i>M. gypseum</i> .
Dogs	5	0.3	
Domestic cats	3	1.0	
Horses	5	0.5	No dermatophytiae linked to the fungi <i>M. canis</i> and <i>T. mentagrophytes</i> after being in direct contact with diseased animals.
Ponies	3	0.3	
Camels	2	5.0	
Bears	2	3.0	
Leopards	2	4.0	
Hyenas	2	2.0	No dermatophytiae linked to the fungi <i>M. canis</i> and <i>T. mentagrophytes</i> after being in direct contact with sources of infection.
Servals	2	3.0	
Ocelots	2	2.0	
Lions	2	3.0	
Tigers	3	7.0	
Nasuas	3	0.5	
Civets	2	1.0	
Rabbits	7	1.5	No symptoms of disease after infection with virulent cultures of the fungi <i>T. sarkisovii</i> and <i>M. gypseum</i> .
Dogs	3	0.5	
Domestic cats	3	1.5	
Black panthers	2	5.0	No dermatophytiae linked to the fungi <i>M. canis</i> , <i>T. mentagrophytes</i> and <i>T. verrucosum</i> after being in direct contact with sources of infection.
Tigers	5	7.0	
Geese	6	3.0	
Bears	3	1.0	
Dogs	8	0.5	
Llamas	2	3.0	

Table 10			
Type of animals	Number	Dosage (cm ³)	Effectiveness
Black panthers	5	7.0	Affected by microsporosis linked to the fungi <i>M. canis</i> . Recovery took place 12-25 days after immunization.
Black panthers	3	4.0	
Horses	3	1.0	
Ponies	2	0.5	
Lions	3	10	
Tigers	3	10	
Dogs	4	0.5	
Bear	1	5.0	
Hycna	1	5.0	
Domestic cats	15	1.5	Affected by microsporosis linked to the fungi <i>M. canis</i> . Recovery took place 10-20 days after immunization.
Dogs	5	0.5	
Horses	5	0.7	
Black panther	1	6.0	Affected by trichophytosis linked to the fungi <i>T. mentagrophytes</i> . Recovery took place 12-15 days after immunization.
Red foxes	4	1.0	
Bears	2	5.0	
Mountain sheep	1	7.0	
Horses	15	1.0	Affected by microsporosis linked to the fungi <i>M. equinum</i> . Recovery took place 12-20 days after immunization.

Bibliography

- (1) Aisenberg, A.A., Noskow, A.I., Kolovatsky, P.P. "Primenenie Yuglona v Veterinarii" in Scientific and Technical Information Bulletin of the State and Scientific Control Committee under the Moldavian Council of Ministers (1958), p. 88.

5

- (2) USSR Patent No. 548947 (1976).